Supplementary Data 1. Questionnaire sheet

This is a questionnaire for monitoring local and systemic adverse reactions after COVID-19 vaccination. To understand the adverse reactions after receiving the COVID-19 vaccine, we are going to collect and use personal information for monitoring all respondets as follows. All information will be anonymized and fully encrypted before analysis.

PERSONAL DATA

PER	SONAL DATA		
1	Name	:	
2	Date of birth	:	dd/mm/yy
3	Gender	:	Female
			Male
4	Weight & Height	:	kg/cm
5	Blood type	:	А
			В
			AB
			0
6	Employment /job	:	Healthcare worker (Doctor, Nurse, Pharmacists, etc.)
			Civil servant
			Student
			Private worker
			Unemployed
			Others
VA	CCINATION DATA		
1.	Type of vaccine dose 1	:	Astrazeneca
			Sinopharm
			Moderna
			Novavax
			Pfizer
			Sinovac
			etc
2.	Type of vaccine dose 2	:	Astrazeneca
			Sinopharm
			Moderna
			NT
			Novavax
			Novavax Pfizer Sinovac

				etc
				Not yet vaccinated 2nd dose
3	Type of vaccine dose 3	:		Astrazeneca
				Sinopharm
				Moderna
				Novavax
				Pfizer
				Sinovac
				etc
				Not yet vaccinated 3rd dose
3	Type of vaccine dose 4	:		Astrazeneca
				Sinopharm
				Moderna
				Novavax
				Pfizer
				Sinovac
				etc
				Not yet vaccinated 4th dose
4	Date of last vaccination	:		dd/mm/yy
нıя	STORY OF COVID-19			
1	Covid-19 infection following dose 1 of	:		No
	immunization			Yes
2	Covid-19 infection following dose 2 of immunization	:		No
				Yes
3	Covid-19 infection following dose 3 of	:		No
	immunization			Yes
N / T				
	DICAL CONDITION		_	No
1.	Comorbidity	:		No
				Yes
2	History of allergy	:		No
				Food allergy
				Drug allergy
3	Hospitalization in the last 3 months	:		No
-		•		Yes
4	History of medication in the last 6	:		No
-	- month of mean and the fast of	•		

	months		Yes
AEI	FIS AND THE SEVERITY LEVELS		
1	Pain at injection site	:	Absent
			Does not interfere with activity
			Interferes with activity or repeated use of non-
			narcotic pain reliever >24hrs Prevents daily Activity or repeated Use of narcotic pain Reliever
			Emergency room visit or hospitalization
2	Tenderness/ Soreness	:	Absent
			Mild discomfort to touch
			Discomfort with movement
			Significant discomfort at rest
			Emergency room (ER) visit or hospitalization
3	Redness/ Erythema	:	Absent
			2.5-5cm
			5.1-10cm
			>10 cm
			Necrosis or exfoliative dermatitis
4	Swelling/ induration	:	Absent
			2.5-5cm and does not interfere with daily activity
			5.1-10cm or interferes with daily activity
			>10 cm prevents daily activity
			Necrosis
5	Pruritus associated with injection	:	Absence of any Pruritus
			Itching localized to injection site and relieved spontaneously or With<48 hours treatment Itching beyond injection site not generalized or localized itching Requiring >48 hours treatment
			Itching causing inability to perform usual social & functional activities NA
6	Pain in the limb	:	Absent
0		·	Does not interfere with activity
			Repeated use of nonnarcotic pain reliever >24 hours or
			interferes with daily activity Any use of narcotic pain reliever or prevents daily activity
			Emergency room (ER) visit or hospitalization
7	Fever	:	<38°C (<100.4°F)
			38.0-38.4ºC(100.4101.1ºF)
			38.5-38.9º C (101.2- 102.0ºF)
			39.0-40°C (102.1-104°F)

			> 40°C (>104°F)
8	Nausea/ Vomiting	:	Absent
			No interference with daily activity or 1-2 episodes/24 hours
			Some interference with daily activity or > 2 episodes/24 hours
			Prevents daily activity, requires outpatient IV hydration
9	Headache	:	Emergency room (ER) visit or hospitalization for hypotensive shock Absent
,	Treaderic	•	No interference with daily activity
			Some interference with daily activity or Repeated use of
			nonnarcotic pain reliever Significant, prevents daily Activity or repeated Use of
			narcotic pain Reliever Emergency room (ER) visit or hospitalization
10	Fatigue	:	Absent
	-		No interference with daily activity
			Some interference with daily activity
			Significant, prevents daily activity
			Emergency room (ER) visit or hospitalization
11	Myalgia	:	Absent
			No interference with daily activity
			Some interference with daily activity
			Significant, prevents daily activity
			Emergency room (ER) visit or hospitalization
12	Acute Allergic Reaction	:	Absent
			No interference with daily activity
			Some interference with daily activity
			Prevents daily activity
			Emergency room (ER) visit or hospitalization
13	Rash	:	Rash Absent
			Rashes covering <10% BSA with or without symptoms
			(pruritus, burning, tightness) Rashes covering 10-30 %BSA (Body Surface Area) with or without symptoms (pruritus, burning, tightness),
			Interferes with daily activity Rashes covering >30 %BSA with or without symptoms (pruritus, burning, tightness), prevents With daily
			activity NA
14	Joint pain	:	Absent
	-		Does not interfere with daily activity
			Repeated use of nonnarcotic pain reliever > 24 hours or
			interferes with daily activity Any use of narcotic pain reliever or prevents daily
			activity Emergency room (ER) visit or hospitalization

- 15 Any other Systemic AE's
- : 🗋 Absent

: 🗆 _

- Does not interfere with daily activity
- □ Interferes with daily activity
- Prevents daily activity
- □ Emergency room (ER) visit or hospitalization
- 16 Has any medication/therapy been given or taken to reduce the side effects of existing vaccinations?

Supplementary Data 2. Grading Scales for Local Adverse Events and Systemic

No.	Event Name	None	Mild	Moderate	Severe	Potentially life threatening
1	Pain at injection site	Absent	Does not interfere with activity	Interferes with activity or repeated use of non- narcotic pain reliever >24hrs	Prevents daily Activity or repeated Use of narcotic pain Reliever	Emergency room visit or hospitalization
2	Tenderness/ Soreness	Absent	Mild discomfort to touch	Discomfort with movement	Significant discomfort at rest	Emergency room (ER) visit or hospitalization
3	Redness/ Erythema	Absent	2.5-5cm	5.1-10cm	>10 cm	Necrosis or exfoliative dermatitis
4	Swelling/ induration	Absent	2.5-5cm and does not interfere with daily activity	5.1-10cm or interferes with daily activity	>10 cm prevents daily activity	Necrosis
5	Pruritus associated with injection	Absence of any Pruritus	Itching localized to injection site and relieved spontaneously or With<48 hours treatment	Itching beyond injection site not generalized or localized itching Requiring >48 hours treatment	Itching causing inability to perform usual social & functional activities	NA
6	Pain in the limb	Absent	Does not interfere with activity	Repeated use of nonnarcotic pain reliever >24 hours or interferes with daily activity	Any use of narcotic pain reliever or prevents daily activity	Emergency room (ER) visit or hospitalization
7	Fever	<38ºC (<100.4ºF)	38.0- 38.4ºC(100.4101.1ºF)	38.5-38.9º C (101.2- 102.0ºF)	39.0-40°C (102.1- 104°F)	> 40°C (>104°F)
8	Nausea/ Vomiting	Absent	No interference with daily activity or 1-2 episodes/24 hours	Some interference with daily activity or > 2 episodes/24 hours	Prevents daily activity, requires outpatient IV hydration	Emergency room (ER) visit or hospitalization for hypotensive shock

9	Headache	Absent	No interference with daily activity	Some interference with daily activity or Repeated use of nonnarcotic pain reliever	Significant, prevents daily Activity or repeated Use of narcotic pain Reliever	Emergency room (ER) visit or hospitalization
10	Fatigue	Absent	No interference with daily activity	Some interference with daily activity	Significant, prevents daily activity	Emergency room (ER) visit or hospitalization
11	Myalgia	Absent	No interference with daily activity	Some interference with daily activity	Significant, prevents daily activity	Emergency room (ER) visit or hospitalization
12	Acute Allergic Reaction	Absent	No interference with daily activity	Some interference with daily activity	Prevents daily activity	Emergency room (ER) visit or hospitalization
13	Rash	Rash Absent	Rashes covering <10% BSA with or without symptoms (pruritus, burning, tightness)	Rashes covering 10- 30 %BSA (Body Surface Area) with or without symptoms (pruritus, burning, tightness), Interferes with daily activity	Rashes covering >30 %BSA with or without symptoms (pruritus, burning, tightness), prevents With daily activity	NA
14	Joint pain	Absent	Does not interfere with daily activity	Repeated use of nonnarcotic pain reliever > 24 hours or interferes with daily activity	Any use of narcotic pain reliever or prevents daily activity	Emergency room (ER) visit or hospitalization
15	Any other Systemic AE's	Absent	Does not interfere with daily activity	Interferes with daily activity	Prevents daily activity	Emergency room (ER) visit or hospitalization